

Durham Research Online

Deposited in DRO:

06 December 2016

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Smith, Martin R. and Hughes, Gareth M. G. and Vargas, María C. and de La Parra, Felipe (2016) 'Sclerites and possible mouthparts of Wiwaxia from the temperate palaeolatitudes of Colombia, South America.', *Lethaia*, 49 (3). pp. 393-397.

Further information on publisher's website:

<http://dx.doi.org/10.1111/let.12154>

Publisher's copyright statement:

This is the accepted version of the following article: Smith, Martin R. and Hughes, Gareth M. G. and Vargas, María C. and de La Parra, Felipe (2016) 'Sclerites and possible mouthparts of Wiwaxia from the temperate palaeolatitudes of Colombia, South America.', *Lethaia*, 49 (3). pp. 393-397, which has been published in final form at <http://dx.doi.org/10.1111/let.12154>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.



Sclerites and possible mouthparts of *Wiwaxia* from the temperate palaeolatitudes of Colombia, South America

Journal:	<i>Lethaia</i>
Manuscript ID:	LET-OA-02-15-0605.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Smith, Martin; Royal Ontario Museum, Palaeobiology Hughes, Gareth; Petrostrat Ltd., Vargas, María; Instituto Colombiano del Petróleo-Ecopetrol, Biostratigraphy team de la Parra, Felipe; Instituto Colombiano del Petróleo-Ecopetrol, Biostratigraphy team
Keywords:	Small Carbonaceous Fossils, <i>Wiwaxia</i> , Cambrian, South America, palynological processing, biogeography

SCHOLARONE™
Manuscripts

View

Sclerites and possible mouthparts of *Wiwaxia* from the temperate
palaeolatitudes of Colombia, South America

Martin R. Smith [ms609@cam.ac.uk], Department of Earth Sciences, Downing Site,
University of Cambridge, Cambridge, CB2 3EQ, UK*

Gareth M. G. Hughes [gareth.hughes@petrostrat.com], PetroStrat Ltd., Tan-Y-Graig,
Parc Caer Seion, Conwy, LL32 8FA, UK*

*M. Carolina Vargas & Felipe de la Parra, Biostratigraphy team, Instituto Colombiano
del Petróleo-Ecopetrol, AA 4185, Bucaramanga, Colombia*

** These authors contributed equally to the current study*

Abstract

The problematic mollusc *Wiwaxia* is perhaps the most widely distributed
non-mineralized Cambrian metazoan, but has only been reported from palaeotropical
latitudes. Here we describe mid-Cambrian (Drumian, c. 504 Ma) sclerites and possible
tooth arrays from the northern Llanos Basin, Colombia, recovered from drilled ditch
cuttings by palynological processing – demonstrating that pristine material and
low-manipulation processing are not essential to the recovery of Small Carbonaceous
Fossils. This, the first report of *Wiwaxia* from South America, substantially expands
Wiwaxia's geographic range into the high palaeolatitudes.

1
2
3 Cambrian lagerstätten such as the Burgess Shale are renowned for their exceptionally
4
5 preserved non-mineralized metazoans. Among the most iconic of these animals is
6
7
8 *Wiwaxia*, a problematic slug-like mollusc covered in imbricating scales and spines.
9
10 Originally interpreted as an annelid worm (Walcott 1911; Butterfield 1990), *Wiwaxia* is
11
12 now recognized as a mollusc based on its radula, creeping ventral foot, and
13
14 aculiferan-like scleritome (Conway Morris 1985; Smith 2012, 2014), although its
15
16 precise position within total-group Mollusca remains unclear.
17
18
19

20
21 The anatomy of *Wiwaxia* has been reconstructed from articulated specimens,
22
23 known from four Burgess Shale-type deposits in North America and China (Conway
24
25 Morris 1985; Zhao *et al.* 1994; Yang *et al.* 2014; Conway Morris *et al.* 2015). The
26
27 organism bears four sclerite morphologies (Conway Morris 1985): depending on their
28
29 position, sclerites are asymmetric (dorsal), rounded and symmetrical (upper lateral and
30
31 anterior), elongate and symmetrical (lower lateral) or sickle-shaped (ventral). Mature
32
33 *Wiwaxia* specimens also exhibit elongate dorso-lateral spines.
34
35
36
37

38
39 Sclerites bear a variable number of ribs and express a thickened margin and a
40
41 microvillar construction (Butterfield 1990; Smith 2014). They are constructed from a
42
43 broad oval blade and narrow, hollow root; this distinctive outline means that isolated
44
45 *Wiwaxia* sclerites are readily identified (Conway Morris 1985; Butterfield 1990).
46
47
48

49 Indeed, lone *Wiwaxia* sclerites have been reported on bedding-surfaces (Ivantsov *et*
50
51 *al.* 2005; Fatka *et al.* 2011; Sun *et al.* 2014; Zhao *et al.* 2015), as phosphatic casts
52
53 (Porter 2004), and – most importantly – as Small Carbonaceous Fossils (SCFs)
54
55 (Butterfield & Harvey 2012; Harvey *et al.* 2012; Pedder 2012).
56
57
58
59
60

SCFs are robust carbon films, typically extracted from pristine siliciclastic mudstones and siltstones through the hand-picking of sieved macerates (Butterfield & Harvey 2012). Owing to their small size and taphonomic recalcitrance, they are widely distributed in space and time, and extend the record of *Wiwaxia* across the modern globe (Fig. 1). Nevertheless, *Wiwaxia* has not been reported outside the low palaeolatitudes – possibly reflecting the increased sampling effort applied to low-palaeolatitude continents. Our material from the Llanos Basin of Colombia, South America, was deposited at a latitude between 50–90° S (McKerrow *et al.* 1992; Torsvik & Cocks 2013), providing the first sampling of SCFs from high palaeolatitudes.

Materials and methods

Mudstone samples were obtained from washed and dried ditch cuttings from the Chigüiro-1 well, which was drilled in 1985 in the northern Llanos Basin, north-east Colombia. Ditch cuttings were produced by the rotational grinding action of the drill bit and brought to the surface by the conveyor-belt action of circulating drilling fluid. Samples were collected and bagged at regular intervals; each sample thus represents the depth of rock drilled since the last sample was taken. Over the *Wiwaxia*-yielding interval (8750–9710 ft), each sample typically corresponds to an interval of 20–30 ft (6–10 m) of rock; each quoted sample depth denotes the bottom of the sampled interval.

5–20 g of each sample was prepared either by standard palynological procedures – maceration by successive applications of HCl and HF, followed by heavy mineral separation in LST FastFloat™, sieving through a 10 µm mesh, and strew mounting (56

1
2
3
4 samples in total; 20 from the *Wiwaxia*-bearing interval, of which 12 yielded *Wiwaxia*) –
5
6 or a low-manipulation procedure incorporating gentle dissolution in HF, sieving
7
8 through a 63 µm gauze, and manual mounting onto glass slides by pipette (21 samples
9
10 from the *Wiwaxia*-bearing interval, 4 yielding *Wiwaxia*).
11
12

13
14 Light micrographs from multiple focal planes were combined using *TuFuse*
15
16 (tawbaware.com); backgrounds and extraneous objects have been removed.
17
18 Measurements were recorded digitally from micrographs. SCFs are deposited at the
19
20 Sedgwick Museum of Earth Sciences, Cambridge (CAMSM); comparative macrofossil
21
22 material is accessioned at the Smithsonian Institution National Museum of Natural
23
24 History (NMNH) and the Royal Ontario Museum (ROM).
25
26
27
28
29

30 **Geological setting**

31
32 Co-occurring acritarchs (in particular *Retisphaeridium dichamerum*, *Eliasum llaniscum*,
33
34 *Adara matutina* and *Cristallinium cambriense*), together with the first downhole
35
36 occurrence of the acritarch *Vulcanisphaera lanugo* in the interval below the last
37
38 persistent downhole occurrence of *Wiwaxia* sclerites, confine the range of *Wiwaxia*
39
40 sclerites to within the *Rugasphaera terranova* acritarch zone (Martin & Dean 1988),
41
42 corresponding to the *Ptychagnostus atavus* / *Tomagnostus fissus* trilobite zone and an
43
44 approximate age of 504 Ma.
45
46
47
48
49

50 The frequent-to-superabundant occurrences of *Siphonophycus* spp. (stromatolite
51
52 filaments) throughout the *Wiwaxia*-bearing interval indicate a shallow water
53
54 depositional environment (Butterfield & Chandler 1992), whereas the regional geology
55
56 indicates a distal shelf setting (Torsvik & Cocks 2013).
57
58
59
60

Sclerites

Twenty-five sclerites were recovered: eighteen through palynological preparations and seven through manual extraction, in approximate proportion to the amount of sample processed by each method. Sclerites ranged from 100–500 µm in maximum dimension; neither the typical size nor the sclerite integrity exhibited meaningful variation between the two extraction methods.

Sclerites could be identified as *Wiwaxia* scales based on their construction from a broad, ribbed blade and hollow root, and by the presence of microvillar chambers (Fig. 2). Sclerites can further be designated as: ventro-lateral (two, Fig. 2a, e), based on a siculate habit; lower lateral (five, Fig. 2b, f), based on an elongate aspect and symmetrical shape; upper lateral (four, Fig. 2c, g), based on their rounded profile and 1:1 height:width ratio; and dorsal (four, Fig. 2d, h), based on their marked asymmetry. Ten incomplete sclerites are insufficiently preserved for a confident assignment. Overall, the sclerites' shape and dimensions correspond to those of articulated *Wiwaxia corrugata* juveniles (Fig. 2k), though the sample size is insufficient to attempt species-level taxonomy. Three specimens displayed superficial pustules with an irregular appearance and distribution that conceivably correspond to surface ornament (Fig. 2i–j).

Mouthparts

Sclerites are not the only tough component of *Wiwaxia*: its mouthparts also have a robust carbonaceous construction and could in principle be preserved as SCFs (Smith 2012). The mouthparts were originally understood to represent a series of two to three

1
2
3
4 denticulate bars (Conway Morris 1985), and on this basis various SCFs from the Mount
5
6 Cap formation (mid-Cambrian of Canada) were compared with the rows of teeth in
7
8 *Wiwaxia* (Harvey & Butterfield 2011). However, a more detailed morphological
9
10 interpretation demonstrates that each ‘bar’ in fact represents an array of around two
11
12 dozen shoehorn-shaped teeth arranged symmetrically about a triangular central tooth
13
14 (Smith 2012), undermining the comparison with the Mount Cap SCFs. Despite the
15
16 robust nature of the teeth, therefore, no convincing representatives are available from
17
18 the SCF record.
19
20
21
22

23
24 Our palynological samples contained four SCF elements that correspond to the
25
26 morphology of the *Wiwaxia* radula (Fig. 3). These comprise 100–250 µm long series
27
28 of four to six teeth – equivalent in size to the mouthparts of the smallest articulated
29
30 *Wiwaxia* specimens (Smith 2012; Zhang *et al.* forthcoming). Each tooth has a narrow
31
32 root that tapers into a broad, flat-ended scoop. Teeth are attached by a basal membrane
33
34 (Fig. 3). In the context of the co-occurring *Wiwaxia* sclerites and the absence of other
35
36 metazoan components, the tooth rows’ distinctive morphology identifies them as
37
38 candidate *Wiwaxia* mouthparts.
39
40
41
42
43
44

45 Discussion

46
47
48 *Wiwaxia*-like mouthparts.—*Wiwaxia*-like mouthparts have not been convincingly
49
50 reported as SCFs – despite the robust construction indicated by the thickness of the
51
52 carbon films and (unlike their co-occurring scales) their association with traces of
53
54 phosphorous (Smith 2012). Phosphorous seemingly typifies the most recalcitrant
55
56
57
58
59
60

components of Burgess Shale-type organisms; it is also associated with the tough dorsal armature of *Hallucigenia* and the most robust teeth in *Ottoia*, both of which occur as SCFs (Caron *et al.* 2013; Smith *et al.* in press).

The recovery of probable *Wiwaxia* mouthparts from the Llanos basin confirms this taphonomic potential. The rarity of mouthpart SCFs relative to sclerites may partly reflect the less distinctive morphology of the tooth rows, but it is also likely that sclerites – which are periodically shed into the water column (Smith 2014) – were originally more abundant than mouthparts, which were swallowed when they were moulted (Smith 2014) and presumably disarticulate during digestion.

Distribution of Wiwaxia.—Although early reconstructions placed Colombia at equatorial palaeolatitudes (Scotese & McKerrow 1990), current palaeogeographic maps indicate a location in a cooler climate belt, whether in a polar (McKerrow *et al.* 1992; Scotese *et al.* 1999; Meert & Lieberman 2004) or temperate (Torsvik & Cocks 2013) setting. Our report of *Wiwaxia* from South America thus represents the first data on its distribution at higher latitudes. The equatorial distribution suggested by previous occurrences of *Wiwaxia* (Fig. 1) thus reflects an acquisition bias – higher latitudes are only represented by the poorly sampled terranes of Africa, Antarctica and South America. As such, *Wiwaxia* seems to represent an environmental generalist, able to colonize habitats regardless of ecological, geographical or climatic constraints.

Implications for microfossil recovery.—SCFs are generally recovered through the manual manipulation of pristine mudstones, reflecting the belief that SCFs are too large or too delicate for recovery by traditional palynological techniques (Butterfield &

Harvey 2012). There is only one report of SCF recovery through palynological preparation: copepod mandibles and *Wiwaxia* sclerite fragments from the Cambrian Nolichucky Shale (Pedder 2012; Harvey & Pedder 2013). These specimens are uniformly smaller than 80 μm , around the minimum size that can be recovered by the manual manipulation approach. Corresponding copepod mandibles recovered from the Deadwood Formation by manual manipulation occupy the 100–500 μm size bracket; this seems to suggest that elements in this larger size range disintegrate during centrifugation, prohibiting their recovery by the palynological technique (Harvey & Pedder 2013).

Our study, however, demonstrates that certain SCFs – *Wiwaxia* sclerites and mouthparts – are robust to palynological processing, even in the 100–500 μm size range. If this is representative of SCFs more generally, processing method alone cannot account for the scarcity of recognizable metazoan microfossils in palynological preparations.

Moreover, this report represents the first recovery of SCFs from washed and dried ditch cuttings, a sampling and preparation method that substantially disaggregates the rock; conventionally, samples from outcrop or pristine cores are preferred. The unrecognized potential of low-grade but abundant ditch cuttings for SCF recovery opens the possibility of a systematic high-throughput sampling of Cambrian wellbore material.

Acknowledgements. We thank the two referees for their constructive comments, and Ecopetrol SA for permission to publish the wellbore material. Jean-Bernard Caron,

Peter Fenton, Doug Erwin and Mark Florence provided access to Burgess Shale material, which was funded in part by a Geological Society of America research grant to M.R.S. M.R.S. is supported by Clare College, Cambridge.

Figure 1. Stratigraphic and geographical distribution of *Wiwaxia*. Silhouettes denote whether sites preserve articulated specimens or isolated sclerites. ‘Wiwaxiid’ material from Sinsk (Ivantsov *et al.* 2005) has not been figured or described and its identity is therefore uncertain. A reported occurrence in South Australia (Emu Bay Shale, Porter 2004) is based on unpublished material and is thus not depicted. The palaeogeographic reconstruction is based on true polar wander data (after Torsvik & Cocks 2013). Colour online.

Figure 2. Sclerites of *Wiwaxia* from the Llanos Basin, Colombia. (a–d), sclerites recovered by manual extraction: (a), CAMSX #####1a (–9130’), ventro-lateral sclerite; (b), CAMSX #####2a (–9050’), lower lateral sclerite; (c), CAMSX #####3a (–9100’), upper lateral sclerite; (d) CAMSX #####3b (–9100’), dorsal sclerite. (e–i), sclerites recovered by palynological processing: (e), CAMSX #####4a (–8750’), ventro-lateral sclerite; (f), CAMSX #####5a (–9810’), lower lateral sclerite; (g), CAMSX #####6a (–9710’), upper lateral or anterior sclerite; (h), CAMSX #####7a (–9650’), dorsal sclerite; (i), CAMSX #####4a (–8750’), enlargement of boxed area in e, demonstrating oval pustules. (j), CAMSX #####2b (–9050’), upper lateral sclerite recovered by

1
2
3
4 manual extraction; enlargement of boxed area demonstrates microvillar construction
5
6 and superficial pustules. (k), NMNH 229901, showing position of sclerites on
7
8 articulated juvenile of *Wiwaxia corrugata*. Scale bars = 100 μm except panels marked
9
10 +, 70 μm . Colour online.

11
12
13
14
15
16
17
18 Figure 3. Possible *Wiwaxia* tooth rows from the Llanos Basin, Colombia, recovered
19
20 through palynological processing; (a), CAMSX #####10a (–9140’); (b), CAMSX
21
22 #####8a (–9010’); (c), CAMSX #####9a (–8990’); (d), CAMSX #####11a (–8440’).
23
24
25 Colour online.

26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

References

- Butterfield, N.J. 1990: A reassessment of the enigmatic Burgess Shale fossil *Wiwaxia corrugata* (Matthew) and its relationship to the polychaete *Canadia spinosa* Walcott. *Paleobiology* 16, 287–303.
- Butterfield, N.J. and Chandler, F.W. 1992: Palaeoenvironmental distribution of Proterozoic microfossils, with an example from the Agu Bay Formation, Baffin Island. *Palaeontology* 35, 943–957.
- Butterfield, N.J. and Harvey, T.H.P. 2012: Small Carbonaceous Fossils (SCFs): a new measure of early Paleozoic paleobiology. *Geology* 40, 71–74.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Caron, J.-B., Smith, M.R. and Harvey, T.H.P. 2013: Beyond the Burgess Shale:
Cambrian microfossils track the rise and fall of hallucigeniid lobopodians.
Proceedings of the Royal Society B: Biological Sciences 280, 20131613.

Conway Morris, S. 1985: The Middle Cambrian metazoan *Wiwaxia corrugata*
(Matthew) from the Burgess Shale and *Ogygopsis* Shale, British Columbia,
Canada. *Philosophical Transactions of the Royal Society of London B:*
Biological Sciences 307, 507–582.

Conway Morris, S., Selden, P.A., Gunther, G., Jamison, P.M. and Robison, R.A. 2015:
New records of Burgess Shale-type taxa from the middle Cambrian of Utah.
Journal of Paleontology, in press.

Fatka, O., Kraft, P. and Szabad, M. 2011: Shallow-water occurrence of *Wiwaxia* in
the Middle Cambrian of the Barrandian area (Czech Republic). *Acta*
Palaeontologica Polonica 56, 871–875.

Harvey, T.H.P. and Butterfield, N.J. 2011: Macro- and microfossils of the Mount Cap
Formation (Early and Middle Cambrian, Northwest Territories). *Geoscience*
Canada 38, 165–173.

Harvey, T.H.P. and Pedder, B.E. 2013: Copepod mandible palynomorphs from the
Nolichucky Shale (Cambrian, Tennessee): implications for the taphonomy and
recovery of Small Carbonaceous Fossils. *Palaios* 28, 278–284.

- Harvey, T.H.P., Ortega-Hernández, J., Lin, J.-P., Zhao, Y.-L. and Butterfield, N.J. 2012: Burgess Shale-type microfossils from the middle Cambrian Kaili Formation, Guizhou Province, China. *Acta Palaeontologica Polonica* 57, 423–436.
- Ivantsov, A.Yu., Zhuravlev, A.Yu., Leguta, A.V., Krassilov, V.A., Melnikova, L.M. and Ushatinskaya, G.T. 2005: Palaeoecology of the Early Cambrian Sinsk biota from the Siberian Platform. *Palaeogeography, Palaeoclimatology, Palaeoecology* 220, 69–88.
- Martin, F. and Dean, W.T. 1988: Middle and Upper Cambrian acritarch and trilobite zonation at Manuels River and Random Island, eastern Newfoundland. *Geological Survey of Canada, Ottawa, Bulletins* 381, 1–91.
- McKerrow, W.S., Scotese, C.R. and Brasier, M.D. 1992: Early Cambrian continental reconstructions. *Journal of the Geological Society* 149, 599–606.
- Meert, J.G. and Lieberman, B.S. 2004: A palaeomagnetic and palaeobiogeographical perspective on latest Neoproterozoic and early Cambrian tectonic events. *Journal of the Geological Society* 161, 477–487.
- Pedder, B.E. 2012: The palynology and stratigraphy of the Cambrian Nolichucky Shale and associated formations at Thorn Hill, Tennessee, USA. University of Sheffield, 231 pp.

- Porter, S.M. 2004: Halkieriids in Middle Cambrian phosphatic limestones from Australia. *Journal of Paleontology* 78, 574–590.
- Scotese, C.R., Boucot, A.J. and McKerrow, W.S. 1999: Gondwanan palaeogeography and palaeoclimatology. *Journal of African Earth Sciences* 28, 98–114.
- Scotese, C.R. and McKerrow, W.S. 1990: Revised world maps and introduction. In McKerrow, W.S., Scotese, C.R. (eds.), *Paleozoic Paleogeography and Biogeography*, 1–21. Geological Society of London, Memoir 12.
- Smith, M.R. 2012: Mouthparts of the Burgess Shale fossils *Odontogriphus* and *Wiwaxia*: implications for the ancestral molluscan radula. *Proceedings of the Royal Society B: Biological Sciences* 279, 4287–4295.
- Smith, M.R. 2014: Ontogeny, morphology and taxonomy of the soft-bodied Cambrian ‘mollusc’ *Wiwaxia*. *Palaeontology* 57, 215–229.
- Smith, M.R., Harvey, T.H.P. and Butterfield, N.J. 2015: The macro- and microfossil record of the middle Cambrian priapulid *Ottoia*. *Palaeontology*, in press.
- Sun, H.-J., Zhao, Y.-L., Peng, J. and Yang, Y.-N. 2014: New *Wiwaxia* material from the Tsinghsutung Formation (Cambrian Series 2) of Eastern Guizhou, China. *Geological Magazine* 151, 339–348.
- Torsvik, T.H. and Cocks, L.R.M. 2013: Gondwana from top to base in space and time. *Gondwana Research* 24, 999–1030.


Walcott, C.D. 1911: Cambrian Geology and Paleontology II, no. 5. Middle Cambrian annelids. *Smithsonian Miscellaneous Collections* 57, 109–144.

Yang, J., Smith, M.R., Lan, T., Hou, J. and Zhang, X. 2014: Articulated *Wiwaxia* from the Cambrian Stage 3 Xiaoshiba Lagerstätte. *Scientific Reports* 4, 4643.

Zhang, Z., Smith, M.R. and Shu, D. in review: New reconstruction of the *Wiwaxia* scleritome, with data from Chengjiang juveniles. *Scientific Reports*.

Zhao, F., Smith, M.R., Yin, Z.-J., Zeng, H., Hu, S.-X., Li, G.-X. and Zhu, M.-Y. 2015: First report of *Wiwaxia* from the Cambrian Chengjiang Lagerstätte. *Geological Magazine* 152, 378–382.

Zhao, Y.-L., Qian, Y. and Li, X.S. 1994: *Wiwaxia* from Early-Middle Cambrian Kaili Formation in Taijiang, Guizhou. *Acta Palaeontologica Sinica* 33, 359–366.



Stratigraphic and geographical distribution of *Wiwaxia*. Silhouettes denote whether sites preserve articulated specimens or isolated sclerites. 'Wiwaxiid' material from Sinsk (Ivantsov *et al.* 2005) has not been figured or described and its identity is therefore unclear. A reported occurrence in south Australia (Emu Bay Shale, Fatka *et al.* 2011) is based on unpublished material and is thus not depicted. The palaeogeographic reconstruction is based on true polar wander data (after Torsvik & Cocks 2013).




Figure 2. Sclerites of *Wiwaxia* from the Llanos Basin, Colombia. (a–d), sclerites recovered by manual extraction: (a), CAMSX #####1a (–9130'), ventro-lateral sclerite; (b), CAMSX #####2a (–9050'), lower lateral sclerite; (c), CAMSX #####3a (–9100'), upper lateral sclerite; (d) CAMSX #####3b (–9100'), dorsal sclerite. (e–i), sclerites recovered by palynological processing: (e), CAMSX #####4a (–8750'), ventro-lateral sclerite; (f), CAMSX #####5a (–9810'), lower lateral sclerite; (g), CAMSX #####6a (–9710'), upper lateral or anterior sclerite; (h), CAMSX #####7a (–9650'), dorsal sclerite; (i), CAMSX #####4a (–8750'), enlargement of boxed area in e, demonstrating oval pustules. (j), CAMSX #####2b (–9050'), upper lateral sclerite recovered by manual extraction; enlargement of boxed area demonstrates microvillar construction and superficial pustules. (k), NMNH 229901, showing position of sclerites on articulated juvenile of *Wiwaxia corrugata*. Scale bars = 100 µm except panels marked +, 70 µm.




Figure 3. Possible *Wiwaxia* tooth rows from the Llanos Basin, Colombia, recovered through palynological processing; (a), CAMSX #####10a (-9140'); (b), CAMSX #####8a (-9010'); (c), CAMSX #####9a (-8990'); (d), CAMSX #####11a (-8440').